

**NATIONAL BOARD FOR HIGHER MATHEMATICS**

**Research Scholarships Screening Test**

**Saturday, January 24, 2009**

**Time Allowed: 150 Minutes**

**Maximum Marks: 40**

**Please read, carefully, the instructions on the following page**

## INSTRUCTIONS TO CANDIDATES

- Please ensure that this booklet contains 12 numbered (and printed) pages. The back of each printed page is blank and can be used for rough work.
- There are **five** sections, containing **ten** questions each, entitled Algebra, Analysis, Topology, Applied Mathematics and Miscellaneous. Answer as many questions as possible. The assessment of the paper will be based on the best **four** sections. Each question carries one point and the maximum possible score is **forty**.
- Answer each question, as directed, in the space provided in the **answer booklet**, which is being supplied separately. This question paper is meant to be retained by you and so **do not** answer questions on it.
- In certain questions you are required to pick out the qualifying statement(s) from multiple choices. None of the statements, or one or more than one statement may qualify. Write **none** if none of the statements qualify, or list the labels of *all* the qualifying statements (amongst (a), (b), and (c)).
- Points will be awarded in the above questions only if **all** the correct choices are made. **There will be no partial credit.**
- $\mathbb{N}$  denotes the set of natural numbers,  $\mathbb{Z}$  - the integers,  $\mathbb{Q}$  - the rationals,  $\mathbb{R}$  - the reals and  $\mathbb{C}$  - the field of complex numbers.  $\mathbb{R}^n$  denotes the  $n$ -dimensional Euclidean space, which is assumed to be endowed with its 'usual' topology. The symbol  $\mathbb{Z}_n$  will denote the ring of integers modulo  $n$ . The symbol  $]a, b[$  will stand for the open interval  $\{x \in \mathbb{R} \mid a < x < b\}$  while  $[a, b]$  will stand for the corresponding closed interval;  $[a, b[$  and  $]a, b]$  will stand for the corresponding left-closed-right-open and left-open-right-closed intervals respectively. The symbol  $I$  will denote the identity matrix of appropriate order. The space of continuous real valued functions on an interval  $[a, b]$  is denoted by  $\mathcal{C}[a, b]$  and is endowed with its usual 'sup' norm.

## Section 1: Algebra

**1.1** Pick out the cases where the given subgroup  $H$  is a normal subgroup of the group  $G$ .

(a)  $G$  is the group of all  $2 \times 2$  invertible upper triangular matrices with real entries, under matrix multiplication, and  $H$  is the subgroup of all such matrices  $(a_{ij})$  such that  $a_{11} = 1$ .

(b)  $G$  is the group of all  $2 \times 2$  invertible upper triangular matrices with real entries, under matrix multiplication, and  $H$  is the subgroup of all such matrices  $(a_{ij})$  such that  $a_{11} = a_{22}$ .

(c)  $G$  is the group of all  $n \times n$  invertible matrices with real entries, under matrix multiplication, and  $H$  is the subgroup of such matrices with positive determinant.

**1.2** Let  $GL(n, \mathbb{R})$  denote the group of all invertible  $n \times n$  matrices with real entries, under matrix multiplication, and let  $SL(n, \mathbb{R})$  denote the subgroup of such matrices whose determinant is equal to unity. Identify the quotient group  $GL(n, \mathbb{R})/SL(n, \mathbb{R})$ .

**1.3** Let  $S_n$  denote the symmetric group of permutations of  $n$  symbols. Does  $S_7$  contain an element of order 10? If 'yes', write down an example of such an element.

**1.4** What is the largest possible order of an element in  $S_7$ ?

**1.5** Write down all the units in the ring  $\mathbb{Z}_8$  of all integers modulo 8.

**1.6** Pick out the cases where the given ideal is a maximal ideal.

(a) The ideal  $15\mathbb{Z}$  in  $\mathbb{Z}$ .

(b) The ideal  $\mathcal{I} = \{f : f(0) = 0\}$  in the ring  $\mathcal{C}[0, 1]$  of all continuous real valued functions on the interval  $[0, 1]$ .

(c) The ideal generated by  $x^3 + x + 1$  in the ring of polynomials  $\mathbb{F}_3[x]$ , where  $\mathbb{F}_3$  is the field of three elements.

**1.7** Let  $A$  be a  $2 \times 2$  matrix with complex entries which is non-zero and non-diagonal. Pick out the cases when  $A$  is diagonalizable.

(a)  $A^2 = I$ .

(b)  $A^2 = 0$ .

(c) All eigenvalues of  $A$  are equal to 2.

**1.8** Let  $\mathbf{x}$  and  $\mathbf{y} \in \mathbb{R}^n$  be two non-zero (column) vectors. Let  $\mathbf{y}^T$  denote the transpose of  $\mathbf{y}$ . Let  $A = \mathbf{xy}^T$ , i.e.  $A = (a_{ij})$  where  $a_{ij} = x_i y_j$ . What is the rank of  $A$ ?

**1.9** Let  $\mathbf{x}$  be a non-zero (column) vector in  $\mathbb{R}^n$ . What is the necessary and sufficient condition for the matrix  $A = I - 2\mathbf{x}\mathbf{x}^T$  to be orthogonal?

**1.10** Let  $A$  be an  $n \times n$  matrix with complex entries. Pick out the true statements.

- (a)  $A$  is always similar to an upper-triangular matrix.
- (b)  $A$  is always similar to a diagonal matrix.
- (c)  $A$  is similar to a block diagonal matrix, with each diagonal block of size strictly less than  $n$ , provided  $A$  has at least 2 distinct eigenvalues.

## Section 2: Analysis

2.1 Evaluate:

$$\lim_{n \rightarrow \infty} n \sin(2\pi en!).$$

2.2 Evaluate:

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{1}{\sqrt{4n^2 - k^2}}.$$

2.3 Pick out the convergent series:

(a)

$$\sum_{n=1}^{\infty} \frac{(n+1)^n}{n^{n+\frac{3}{2}}}.$$

(b)

$$1 + \frac{1}{2^2} + \frac{2^2}{3^3} + \frac{3^3}{4^4} + \dots$$

(c)

$$\sum_{n=1}^{\infty} \sqrt{\frac{1+4^n}{1+5^n}}.$$

2.4 Which of the following functions are continuous?

(a)

$$f(x) = x^2 + \frac{x^2}{1+x^2} + \frac{x^2}{(1+x^2)^2} + \dots, \quad x \in \mathbb{R}.$$

(b)

$$f(x) = \sum_{n=1}^{\infty} (-1)^n \frac{\cos nx}{n^{\frac{3}{2}}}, \quad x \in [-\pi, \pi].$$

(c)

$$f(x) = \sum_{n=1}^{\infty} n^2 x^n, \quad x \in \left[-\frac{1}{2}, \frac{1}{2}\right].$$

2.5 Which of the following functions are uniformly continuous?

(a)  $f(x) = \frac{1}{x}$  in  $(0, 1)$ .

(b)  $f(x) = x^2$  in  $\mathbb{R}$ .

(c)  $f(x) = \sin^2 x$  in  $\mathbb{R}$ .

2.6 Pick out the sequences  $\{f_n\}$  which are uniformly convergent.

(a)

$$f_n(x) = nxe^{-nx} \quad \text{on } (0, \infty).$$

(b)

$$f_n(x) = x^n \quad \text{on } [0, 1].$$

(c)

$$f_n(x) = \frac{\sin nx}{\sqrt{n}} \quad \text{on } \mathbb{R}.$$

**2.7** Which of the following functions are Riemann integrable on the interval  $[0, 1]$ ?

(a)

$$f(x) = \lim_{n \rightarrow \infty} \cos^{2n}(24\pi x).$$

(b)

$$f(x) = \begin{cases} 0, & \text{if } x \text{ is rational} \\ 1, & \text{if } x \text{ is irrational.} \end{cases}$$

(c)

$$f(x) = \begin{cases} \cos x, & \text{if } 0 \leq x \leq \frac{1}{2} \\ \sin x, & \text{if } \frac{1}{2} < x \leq 1. \end{cases}$$

**2.8** Let  $z = x + iy$  be a complex number, where  $x$  and  $y \in \mathbb{R}$ , and let  $f(z) = u(x, y) + iv(x, y)$ , where  $u$  and  $v$  are real valued, be an analytic function on  $\mathbb{C}$ . Express  $f'(z)$  in terms of the partial derivatives of  $u$  and  $v$ .

**2.9** Let  $z \in \mathbb{C}$  be as in the previous question. Find the image of the set  $S = \{z : x > 0, 0 < y < 2\}$  under the transformation  $f(z) = iz + 1$ .

**2.10** Find the residue at  $z = 0$  for the function

$$f(z) = \frac{1 + 2z}{z^2 + z^3}.$$

### Section 3: Topology

**3.1** Let  $X$  be a metric space and let  $f : X \rightarrow \mathbb{R}$  be a continuous function. Pick out the true statements.

- (a)  $f$  always maps Cauchy sequences into Cauchy sequences.
- (b) If  $X$  is compact, then  $f$  always maps Cauchy sequences into Cauchy sequences.
- (c) If  $X = \mathbb{R}^n$ , then  $f$  always maps Cauchy sequences into Cauchy sequences.

**3.2** Let  $B$  be the closed ball in  $\mathbb{R}^2$  with centre at the origin and radius unity. Pick out the true statements.

- (a) There exists a continuous function  $f : B \rightarrow \mathbb{R}$  which is one-one.
- (b) There exists a continuous function  $f : B \rightarrow \mathbb{R}$  which is onto.
- (a) There exists a continuous function  $f : B \rightarrow \mathbb{R}$  which is one-one and onto.

**3.3** Let  $A$  and  $B$  be subsets of  $\mathbb{R}$ . Define

$$C = \{a + b : a \in A, b \in B\}.$$

Pick out the true statements.

- (a)  $C$  is closed if  $A$  and  $B$  are closed.
- (b)  $C$  is closed if  $A$  is closed and  $B$  is compact.
- (c)  $C$  is compact if  $A$  is closed and  $B$  is compact.

**3.4** Which of the following subsets of  $\mathbb{R}^2$  are compact?

- (a)  $\{(x, y) : xy = 1\}$
- (b)  $\{(x, y) : x^{\frac{2}{3}} + y^{\frac{2}{3}} = 1\}$
- (c)  $\{(x, y) : x^2 + y^2 < 1\}$

**3.5** Which of the following sets in  $\mathbb{R}^2$  are connected?

- (a)  $\{(x, y) : x^2y^2 = 1\}$
- (b)  $\{(x, y) : x^2 + y^2 = 1\}$
- (c)  $\{(x, y) : 1 < x^2 + y^2 < 2\}$

**3.6** Let  $\mathcal{P}$  denote the set of all polynomials in the real variable  $x$  which varies over the interval  $[0, 1]$ . What is the closure of  $\mathcal{P}$  in  $\mathcal{C}[0, 1]$  (with its usual sup-norm topology)?

**3.7** Let  $\{f_n\}$  be a sequence of functions which are continuous over  $[0, 1]$  and continuously differentiable in  $]0, 1[$ . Assume that  $|f_n(x)| \leq 1$  and that  $|f'_n(x)| \leq 1$  for all  $x \in ]0, 1[$  and for each positive integer  $n$ . Pick out the true statements.

- (a)  $f_n$  is uniformly continuous for each  $n$ .
- (b)  $\{f_n\}$  is a convergent sequence in  $\mathcal{C}[0, 1]$ .
- (c)  $\{f_n\}$  contains a subsequence which converges in  $\mathcal{C}[0, 1]$ .

**3.8** Pick out the true statements.

- (a) Let  $f : [0, 2] \rightarrow [0, 1]$  be a continuous function. Then, there always exists  $x \in [0, 1]$  such that  $f(x) = x$ .
- (b) Let  $f : [0, 1] \rightarrow [0, 1]$  be a continuous function which is continuously differentiable in  $]0, 1[$  and such that  $|f'(x)| \leq \frac{1}{2}$  for all  $x \in ]0, 1[$ . Then, there exists a unique  $x \in [0, 1]$  such that  $f(x) = x$ .
- (c) Let  $S = \{\mathbf{p} = (x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$ . Let  $f : S \rightarrow S$  be a continuous function. Then, there always exists  $\mathbf{p} \in S$  such that  $f(\mathbf{p}) = \mathbf{p}$ .

**3.9** Let  $(X, d)$  be a metric space. Let  $A$  and  $B$  be subsets of  $X$ . Define

$$d(A, B) = \inf\{d(a, b) : a \in A, b \in B\}.$$

For  $x \in X$ , define

$$d(x, A) = \inf\{d(x, a) : a \in A\}.$$

Pick out the true statements.

- (a) The function  $x \mapsto d(x, A)$  is uniformly continuous.
- (b)  $d(x, A) = 0$  if, and only if,  $x \in A$ .
- (c)  $d(A, B) = 0$  implies that  $A \cap B \neq \emptyset$ .

**3.10** Let

$$B = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1\} \text{ and } D = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 < 1\}.$$

Pick out the true statements.

- (a) Given a continuous function  $g : B \rightarrow \mathbb{R}$ , there always exists a continuous function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  such that  $f = g$  on  $B$ .
- (b) Given a continuous function  $g : D \rightarrow \mathbb{R}$ , there always exists a continuous function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  such that  $f = g$  on  $D$ .
- (c) There exists a continuous function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  such that  $f \equiv 1$  on the set  $\{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = \frac{3}{2}\}$  and  $f \equiv 0$  on the set  $B \cup \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \geq 2\}$ .



## Section 4: Applied Mathematics

**4.1** A spherical ball of volatile material evaporates (*i.e.* its volume decreases) at a rate proportional to its surface area. If the initial radius is  $r_0$  and at time  $t = 1$ , the radius is  $r_0/2$ , find the time at which the ball disappears completely.

**4.2** A body of mass  $m$  falling from rest under gravity experiences air resistance proportional to the square of its velocity. Write down the initial value problem for the vertical displacement  $x$  of the body.

**4.3** A body falling from rest under gravity travels a distance  $y$  and has a velocity  $v$  at time  $t$ . Write down the relationship between  $v$  and  $y$ .

**4.4** Let  $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$  and let  $\mathbf{v}(\mathbf{x}) = \mathbf{x}$ . Apply Gauss' divergence theorem to  $\mathbf{v}$  over the unit ball

$$B = \{\mathbf{x} \in \mathbb{R}^n : \sum_{i=1}^n x_i^2 \leq 1\}$$

and deduce the relationship between  $\omega_n$ , the ( $n$ -dimensional) volume of  $B$ , and  $\sigma_n$ , the  $((n - 1)$ -dimensional) surface measure of  $B$ .

**4.5** Write down the general solution of the linear system:

$$\begin{aligned} \frac{dx}{dt} &= x + y \\ \frac{dy}{dt} &= 4x - 2y. \end{aligned}$$

**4.6** What is the smallest positive value of  $\lambda$  such that the problem:

$$\begin{aligned} u'' + \lambda u &= 0 \text{ in } ]0, 1[ \\ u(0) = u(1) &\text{ and } u'(0) = u'(1) \end{aligned}$$

has a solution such that  $u \not\equiv 0$ ?

**4.7** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function. Write down the solution of the problem:

$$\begin{aligned} \frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 u}{\partial x^2} &= 0, \quad t > 0, \\ u(x, 0) &= f(x), \quad x \in \mathbb{R}, \\ \frac{\partial u}{\partial t}(x, 0) &= 0, \quad x \in \mathbb{R}. \end{aligned}$$

**4.8** Use duality to find the optimal value of the cost function in the following linear programming problem:

$$\begin{aligned} \text{Max. } &x + y + z \\ \text{such that } &3x + 2y + 2z = 1, \\ &x \geq 0, \quad y \geq 0, \quad z \geq 0. \end{aligned}$$

**4.9** The value of  $\sqrt{10}$  is computed by solving the equation  $x^2 = 10$  using the Newton-Raphson method. Starting from some value  $x_0 > 0$ , write down the iteration scheme.

**4.10** Write down the Laplace transform  $F(s)$  of the function  $f(x) = x^3$ .

## Section 5: Miscellaneous

**5.1** Let  $P = (0, 1)$  and  $Q = (4, 1)$  be points on the plane. Let  $A$  be a point which moves on the  $x$ -axis between the points  $(0, 0)$  and  $(4, 0)$ . Let  $B$  be a point which moves on the line  $y = 2$  between the points  $(0, 2)$  and  $(4, 2)$ . Consider all possible paths consisting of the line segments  $PA$ ,  $AB$  and  $BQ$ . What is the shortest possible length of such a path?

**5.2** A convex polygon has its interior angles in arithmetic progression, the least angle being  $120^\circ$  and the common difference being  $5^\circ$ . Find the number of sides of the polygon.

**5.3** Let  $a$ ,  $b$  and  $c$  be the lengths of the sides of an arbitrary triangle. Define

$$x = \frac{ab + bc + ca}{a^2 + b^2 + c^2}.$$

Pick out the true statements.

- (a)  $\frac{1}{2} \leq x \leq 2$ .
- (b)  $\frac{1}{2} \leq x \leq 1$ .
- (c)  $\frac{1}{2} < x \leq 1$ .

**5.4** What is the maximum number of pieces that can be obtained from a pizza by making 7 cuts with a knife?

**5.5** In arithmetic base 3, a number is expressed as 210100. Find its square root and express it in base 3.

**5.6** Evaluate:

$$\left( \frac{-1 + i\sqrt{3}}{\sqrt{2} + i\sqrt{2}} \right)^{20}.$$

**5.7** Let  $n$  be a fixed positive integer and let  $C_r$  stand for the usual binomial coefficients *i.e.*, the number of ways of choosing  $r$  objects from  $n$  objects. Evaluate:

$$C_1 + 2C_2 + \cdots + nC_n.$$

**5.8** Let  $x$ ,  $y$  and  $z$  be real numbers such that  $x^2 + y^2 + z^2 = 1$ . Find the maximum and minimum values of  $2x + 3y + z$ .

**5.9** Let  $x_0 = 0$ . For  $n \geq 0$ , define

$$x_{n+1} = x_n^2 + \frac{1}{4}.$$

Pick out the true statements:

- (a) The sequence  $\{x_n\}$  is bounded.
- (b) The sequence  $\{x_n\}$  is monotonic.
- (a) The sequence  $\{x_n\}$  is convergent.

**5.10** Seven tickets are numbered consecutively from 1 to 7. Two of them are selected in order without replacement. Let  $A$  denote the event that the numbers on the two tickets add up to 9. Let  $B$  be the event that the numbers on the two tickets differ by 3. If each draw has equal probability  $\frac{1}{42}$  (the draw  $(1, 7)$  being considered as distinct from the draw  $(7, 1)$ , for example) find the probability  $P(B|A)$ .